


1/10

Fig. 1

BEST AVAILABLE COPY




2/10

Fig. 2


Human hepatocytes (12 years old) used as an antigen
to prepare a monoclonal antibody

Primary culture

At 30 days




At 43 days



Subculture (3 passages) → immunizing mice

After 33 days



Bar, 100 μ m


3/10

Fig.3



Human tissue (62 years old)

Hybridoma culture supernatant (K8223)

K8223



No primary antibody




Bar, 50 μ m

PP: Portal region

4/10

Fig. 4




Hybridoma culture supernatant: No.23
Isolated human hepatocytes (49 years old)


5/10

Fig. 5

At day 8 in culture



At day 1 in culture




Before sorting

Bar, 100 μ m


Hybridoma culture supernatant: No.23
Isolated human hepatocytes (49 years old)

At day 1 in culture



R2 fraction

At day 8 in culture




Bar, 100 μ m

5/1/10

Fig. 5/1


Hybridoma culture supernatant: No.23
Isolated human hepatocytes (49 years old)

At day 1 in culture



R3 fraction

At day 8 in culture



Bar, 100 μ m


5/2/10

Fig. 5/2

Cultured human hepatocytes
(12 years old, male, 5 passages of subculture)

6/10

Fig. 6






7/10

Fig. 7

Spheroid formation of cultured hepatocytes on MATRIGEL

Monolayer culture





9/10

Fig. 9




Fig. 10

